

Automorphisms Inner in the Local Multiplier Algebra and Connes Spectrum

Raluca Dumitru, Costel Peligrad and Bogdan Visinescu

ABSTRACT: We prove that if (A, G, α) is a C^* -dynamical system with G abelian and if g_0 is in the inner centre of G then $g_0 \in \Gamma_B(\alpha)^\perp$. Conversely if G is compact and $Sp(\alpha)/\Gamma(\alpha)$ is finite, we prove that if $g_0 \in \Gamma_B(\alpha)^\perp$ then g_0 is in the inner centre of G (Proposition 3). We then prove that for A prime and G compact, not necessarily abelian, the first part of Proposition 3 can be extended to the case when α_{g_0} is inner in $M_{loc}(A)^\alpha$ (Proposition 7).

KEYWORDS: C^* -algebras, automorphism groups.

AMS SUBJECT CLASSIFICATION: Primary 46L55, Secondary 46L40

In [8] and [4], the C^* -algebra of local multipliers was used for the first time, under the name of algebra of essential (or limit) multipliers. In [3], this algebra is extensively used in the problem of "strong primeness" of fixed point algebras of C^* -dynamical systems with compact abelian groups. The recent book of P.Ara and M. Mathieu [1] contains a systematic study of the algebra of local multipliers, $M_{loc}(A)$ and its applications.

$M_{loc}(A)$ is by definition the inductive limit of the multiplier algebras $M(I)$, where $I \subset A$ is an essential ideal of A . If $I \subset J$ are two such ideals and z_I, z_J are the open central projections in A'' corresponding to I , respectively J then the mapping $x \xrightarrow{\varphi_{JI}} x \cdot z_I$ is an injective morphism of $M(J)$ into $M(I)$. $M_{loc}(A)$ should be understood as the inductive limit of $(M(I), \varphi_{JI})$.

In this paper we will obtain some results that relate the Connes spectrum with the innerness of automorphisms in the fixed point algebra $M_{loc}(A)^\alpha$.

The results are variants and/or extensions of corresponding results of Olesen, Pedersen and Stormer [6], [7], [9] and Ara and Mathieu [1] to the case of local multiplier algebras.

For definitions and basic results on Arveson spectrum, Connes spectrum and Borchers spectrum we send to [9].

In particular, if (A, G, α) is a C^* -dynamical system with G compact, abelian group and $\gamma \in \hat{G}$ we denote

$$A_\gamma = \{x \in A \mid \alpha_t(x) = \langle t, \gamma \rangle x, \forall t \in G\}$$

For $\gamma = e$, the unit of G , we set $A_e = A^\alpha$.

Then the Arveson spectrum is defined by:

$$Sp(\alpha) = \{\gamma \in \hat{G} \mid A_\gamma \neq (0)\}.$$

$\mathcal{H}^\alpha(A)$ denotes the set of all α -invariant, hereditary C^* -subalgebras of A and $\mathcal{H}_B^\alpha(A)$ the set of all α -invariant, hereditary subalgebras of A ($C \in \mathcal{H}_B^\alpha(A)$ if $C \in \mathcal{H}^\alpha(A)$ and the ideal $\overline{ACA}$ is essential in A). Define the Connes spectrum and the Borchers spectrum as follows:

$$\Gamma(\alpha) = \bigcap_{C \in \mathcal{H}^\alpha(A)} Sp(\alpha|_C)$$

$$\Gamma_B(\alpha) = \bigcap_{C \in \mathcal{H}_B^\alpha(A)} Sp(\alpha|_C)$$

Then $\Gamma(\alpha) \subseteq \Gamma_B(\alpha) \subseteq Sp(\alpha)$. $\Gamma(\alpha)$ is a subgroup of $\hat{G}$.

The corresponding spectra for actions of compact, non-abelian groups are defined in [5], [10].

Definition 1. If (A, G, α) is a C^* -dynamical system we define $H \subset G$, the inner centre of G , as the set of all $t \in G$ for which there is an essential, α -invariant ideal $I \subset A$ such that $\alpha_t = \text{Ad } u$ for some $u \in M(I)^\alpha$.

Remark 2. Since $M(I)^\alpha \subset M_{loc}(A)^\alpha$ for every essential ideal $I \subset A$, it is clear that every automorphism α_t with $t \in H$ is inner in $M_{loc}(A)^\alpha$. Here $M_{loc}(A)^\alpha$ denotes the fixed point algebra of $M_{loc}(A)$ under the extensions of all α_t , $t \in G$.

Our first result is concerned with automorphisms in the inner centre of G .

Proposition 3. Let (A, G, α) be a C^* -dynamical system with G locally compact abelian. If $t \in H$ (the inner centre of G), then $\langle t, \sigma \rangle = 1$ for every $\sigma \in \Gamma_B(\alpha)$. The converse holds if G is compact and $Sp(\alpha)/\Gamma(\alpha)$ is finite.

Proof. The proof of the direct implication is a modification of the corresponding result of Olesen ([9] Theorem 8.9.7) for multiplier algebras.

Let $\alpha_t = \text{Ad } u$, $u \in M(I)^\alpha$ for some essential, α -invariant ideal $I \subset A$. Then, since $\Gamma_B(\alpha) = \Gamma_B(\alpha|_I)$ the proof of the first part of ([9] theorem 8.9.7) carries over word for word with A replaced by I .

Conversely, let G be a compact group and $Sp(\alpha)/\Gamma(\alpha)$ finite. We mention that we do not require A to be simple as in ([9] Theorem 8.9.7).

Let $t_0 \in G$ be such that $\langle t_0, \sigma \rangle = 1$, $\forall \sigma \in \Gamma_B(\alpha)$. Then, by ([9] Proposition 8.8.7) there exists $C \in \mathcal{H}_B^\alpha(A)$ such that $Sp(\alpha|_C) \subseteq \Gamma_B(\alpha) + \Gamma(\alpha)$.

Let $\sigma \in Sp(\alpha|_C)$. Then, there are $\sigma_1 \in \Gamma_B(\alpha)$ and $\sigma_2 \in \Gamma(\alpha) \subseteq \Gamma_B(\alpha)$, such that $\sigma = \sigma_1 + \sigma_2$. Then since t_0 annihilates $\Gamma_B(\alpha)$, it follows that $\langle t_0, \sigma \rangle = 1$. Hence $\alpha_{t_0}|_C = \text{id}$. Let $I = \overline{ACA}$ be the two sided ideal generated by C . Since $C \in \mathcal{H}_B^\alpha(A)$, I is an essential, α -invariant ideal of A . Let $p \in C''' \subset I''$ be the open projection associated to C . Then, the central support $c(p)$ in I'' equals 1. By ([9], Lemma 8.9.1) applied to I'' with $w = p$ we get a unique unitary $u \in I''$ such that $up = pu = p$ and $\alpha_{t_0}''|_{I''} = \text{Ad } u$ and hence $\alpha_{t_0}|_I = \text{Ad } u$.

Since G is abelian, $\alpha_t''(u)$ satisfies all the properties of u . By uniqueness of u with the above properties we get $\alpha_t''(u) = u$ for every $t \in G$.

Define $I_0 = \{x \in I \mid ux \in I\}$. Then I_0 is a two sided, closed ideal of I . Indeed, I_0 is clearly a right ideal. It is also a left ideal since for $y \in I$ we have $uyx = uyu^*ux = \alpha_{t_0}(y)ux \in I$. Since p is the open projection associated with C , we have $px = x$ for every $x \in C$. Therefore, if $x \in C$, $ux = upx = px = x \in C \subset I$. Thus, $C \subseteq I_0$. Since the ideal generated by C is I , it follows that $I_0 = I$ and therefore u is a left multiplier of I . Similarly, one can show that u is a right multiplier of I and so $u \in M(I)$. The proof is complete. $\square$

Corollary 4. *If (A, G, α) is a C^* -dynamical system with G compact abelian group and $Sp(\alpha)$ finite, then $\Gamma_B(\alpha)^\perp$ (the annihilator of $\Gamma_B(\alpha)$) coincides with the inner centre. In particular, if A is G -prime, then $\Gamma(\alpha)^\perp (= \Gamma_B(\alpha)^\perp)$ coincides with the inner centre.*

This corollary generalizes ([1], Theorem 4.5.7 and the first part of Theorem 4.5.8).

Our next result extends and improves on some results in [7], ([9] Theorem 8.10.12).

Proposition 5. *Let (A, G, α) be a C^* -dynamical system with G compact, abelian and A prime. Suppose that $Sp(\alpha)/\Gamma(\alpha)$ is finite. The following conditions are equivalent :*

- (i) $Sp(\alpha) = \Gamma(\alpha)$
- (ii) A^α is prime
- (iii) $M_{loc}(A)^\alpha$ is prime
- (iv) The centre of $M_{loc}(A)^\alpha$ is the scalars
- (v) No $\alpha_t \neq \iota$ is implemented by a unitary in $M_{loc}(A)^\alpha$
- (vi) No $\alpha_t \neq \iota$ is in the inner centre of G .

Proof. (i) $\iff$ (ii) by ([9] Theorem 8.10.4)

(ii) $\implies$ (iii) : Since A^α is prime, every ideal of A^α is a prime C^* -algebra. In particular, for every α -invariant ideal $I \subset A$, I^α is prime. Therefore, the multiplier algebra $M(I^\alpha)$ is prime. By ([3] Proposition 3.3), $M_{loc}(A)^\alpha = \lim_{I \text{ invariant ideal}} M(I^\alpha)$. Since $M(I^\alpha)$ are prime, their inductive limit, $M_{loc}(A)^\alpha$ is prime (to show that inductive limits of prime algebras are prime, one can use [1], Lemma 1.2.32).

(iii) $\implies$ (iv) : This is trivial.

(iv) $\implies$ (v) : If $\alpha_{t_0} = Ad \ u$ for $u \in M_{loc}(A)^\alpha$, then for $a \in M_{loc}(A)^\alpha$, $\alpha_{t_0}(a) = a = uau^*$. Hence u belongs to the centre of $M_{loc}(A)^\alpha$. Therefore $u = \lambda \cdot 1$ for some scalar and $\alpha_{t_0} = \iota$.

(v) $\implies$ (vi) : Evident since the inner centre is included in $M_{loc}(A)^\alpha$.

(vi) $\implies$ (i) : If $Sp(\alpha) \not\supseteq \Gamma(\alpha)$ then since $\Gamma(\alpha)$ is a subgroup of $\hat{G}$, there exists $t_0 \in \Gamma(\alpha)^\perp$ but $t_0 \notin Sp(\alpha)^\perp$. By Proposition 3, t_0 belongs to the inner centre of G i.e. there exists an α -invariant ideal I and a unitary $u \in M(I)^\alpha = M(I^\alpha)$ such that $\alpha_{t_0} = Ad \ u$. Therefore $\alpha_{t_0} = id$. Hence $t_0 \in Sp(\alpha)^\perp$, contradiction. $\square$

Our next result is a stronger version of the first part of Proposition 3 for the case of a prime C^* -algebra A and a compact group G . Namely, we will prove that if $\alpha_{t_0} = Ad \ u$, $u \in M_{loc}(A)^\alpha$ (rather than $u \in M(I)^\alpha$ for some essential, α -invariant ideal) then $t_0 \in \Gamma(\alpha)^\perp (= \Gamma_B(\alpha)^\perp$ since A is now prime).

We will prove this for a compact, not necessarily abelian group G . The spectra $Sp(\alpha)$, $\Gamma(\alpha)$, $\tilde{\Gamma}(\alpha)$ for actions of compact, non abelian groups were defined in ([10], [5]). We will define the weak Connes spectrum $\Gamma_\omega(\alpha)$. Let (A, G, α) be a C^* -dynamical system with G compact. Let $\hat{G}$ be the dual of G (i.e. the set of unitary equivalence classes of irreducible representations of G).

For $\pi \in \hat{G}$, $\pi = [\pi_{ij}]$, $i, j = 1, 2, \dots, d_\pi$ (d_π is the dimension of π) let $\chi_\pi(g) = \sum_i \pi_{ii}(g)$ be the character of π .

If $a \in A$, set $P(\pi)(a) = \int_G d_\pi \overline{\chi_\pi(g)} \alpha_g(a) dg$.

Denote

$$A_\pi = \left\{ \int_G d_\pi \overline{\chi_\pi(g)} \alpha_g(a) dg \mid a \in A \right\}$$

Define

$$sp(\alpha) = \{ \pi \in \hat{G} \mid A_\pi \neq (0) \}$$

The Connes spectrum $\Gamma(\alpha)$ and the strong Connes spectrum $\tilde{\Gamma}(\alpha)$ were defined in [5] and used in problems concerning the simplicity and primeness of the crossed products.

We now define the weak Connes spectrum:

$$\Gamma_\omega(\alpha) = \bigcap_{B \in \mathcal{H}^\alpha(A)} sp(\alpha|_B).$$

We have $\tilde{\Gamma}(\alpha) \subseteq \Gamma(\alpha) \subseteq \Gamma_\omega(\alpha)$.

The first inclusion is strict even in the abelian case. The second inclusion is an equality for abelian groups but is strict for non-abelian groups (this is shown by the example 3.9 and Theorem 3.8 in [10] together with Theorem 2.5 in [5]).

Before starting our result, let us make some notations:

For $\pi = [\pi_{ij}] \in \hat{G}$, denote

$$P_{ij}(\pi)(a) = \int d_\pi \overline{\pi_{ji}(g)} \alpha_g(a) dg, \quad a \in A.$$

Let $A_{ij}(\pi) = \{ P_{ij}(\pi)(a) \mid a \in A \}$.

We collect some facts about the bounded, linear mappings $P(\pi)$, $P_{ij}(\pi)$ from A into A in the following:

- Lemma 6.** (i) $P(\pi) = \sum_{i=1, d_\pi} P_{ii}(\pi)$
(ii) $P_{ij}(\pi)P_{kl}(\pi) = \delta_{il}P_{kj}(\pi)$, where δ_{il} is the Kronecker symbol.
(iii) $P(\pi)P_{ij}(\pi) = P_{ij}(\pi) = P_{ij}(\pi)P(\pi)$
(iv) $P(\pi)^2 = P(\pi)$
(v) $\alpha_g(P_{ij}(\pi)(a)) = \sum_{k=1, d_\pi} P_{ik}(\pi)(a) \cdot \pi_{kj}(g)$
(vi) If $P_{i_0 j_0}(\pi)(a) \neq 0$, then $P_{i_0 k}(\pi)(a) \neq 0$ for every $k = 1, 2, \dots, d_\pi$
and $\{P_{i_0 k}(\pi)(a)\}_{k=1}^{d_\pi}$ are linearly independent.
(vii) If $A_\pi \neq (0)$ then $A_{ij}(\pi) \neq 0$ for every $i, j = 1, 2, \dots, d_\pi$.
(viii) The subspaces $A_\pi \subset A$ are α -invariant.

Proof. (i) follows trivially from definitions.

(ii), (iii), (iv) follow from $\pi(gg') = \pi(g)\pi(g')$ (π is a representation of G) and from Weil's orthogonality relations ([2], 2.2.39).

(v) is easily seen since π is a representation, $\pi(gg') = \pi(g)\pi(g')$ ([2], 2.2.39).

(vi) If $P_{i_0 k}(\pi)(a) = 0$, then by (ii), $P_{k j_0}(\pi)(a)P_{i_0 k}(\pi)(a) = P_{i_0 j_0}(\pi)(a) = 0$, contradiction. So $P_{i_0 k}(\pi)(a) \neq 0$, $k = 1, 2, \dots, d_\pi$ if $P_{i_0 j_0}(\pi)(a) \neq 0$. Let now $\sum_k \lambda_k P_{i_0 k}(\pi)(a) = 0$. Applying $P_{k_0 j_0}(\pi)$ to the previous equality, it follows that $\lambda_{k_0} P_{i_0 j_0}(\pi)(a) = 0$ and hence $\lambda_{k_0} = 0$ for every $k_0 = 1, 2, \dots, d_\pi$.

(vii) If $A_\pi \neq 0$, then there is $a \in A$ such that $P(\pi)(a) = \sum P_{ii}(\pi)(a) \neq 0$. Hence $P_{i_0 i_0}(\pi)(a) \neq 0$ for some i_0 .

By (vi) $P_{i_0 k}(\pi)(a) \neq 0$ for every $k = 1, 2, \dots, d_\pi$.

Let $b = P_{i_0 i}(\pi)(a)$. Then $b \neq 0$ and $P_{ij}(\pi)(b) = P_{i_0 j}(\pi)(a) \neq 0$. So $P_{ij}(\pi)(b) \in A_{ij}(\pi) \neq (0)$.

(viii) Follows from (ii), (iii), (iv), and (v). $\square$

We can now prove:

Proposition 7. Let (A, G, α) be a C^* -dynamical system with G compact and A prime. If $\alpha_{g_0} = \text{Ad } u_0$ for some $u_0 \in M_{\text{loc}}(A)^\alpha$, then $g_0 \in \Gamma_\omega(\alpha)^\perp (\subseteq \Gamma(\alpha)^\perp \subseteq \tilde{\Gamma}(\alpha)^\perp)$.

Proof. Let $\epsilon > 0$. By ([3] Proposition 3.3 and [11] Proposition L.2.2) there exists an α -invariant ideal $I \subset A$ and an unitary $u \in M(I)^\alpha$ such that $\|u_0 - u\| < \frac{\epsilon}{4}$. Let $\lambda_0 \in sp(u)$ and f be a positive continuous function on the unit circle $\mathbb{T}$ with $\text{supp } f \subseteq \{\lambda \in \mathbb{T} \mid |\lambda - \lambda_0| < \frac{\epsilon}{4}\}$. Let also h be a positive, continuous function on the unit circle such that $0 \leq h \leq 1$, $h|_{\text{supp } f} \equiv 1$ and $h(\lambda) = 0$ for every $\lambda \in \mathbb{T}$ with

$|\lambda - \lambda_0| \geq \frac{\epsilon}{4}$. Then $f(u), h(u) \in M(I)^\alpha$ and $f(u)h(u) = f(u)$. Let $C = \overline{f(u)If(u)}$ (norm closure). Then $C \in \mathcal{H}^\alpha(I) \subseteq \mathcal{H}^\alpha(A)$.
Let $x \in C$. Then:

$$\begin{aligned} \|\alpha_{g_0}(x) - x\| &= \|u_0xu_0^* - x\| = \|u_0x - xu_0\| \\ &\leq \|(u_0 - u)x - x(u_0 - u)\| + \|ux - xu\| \\ &< \frac{\epsilon}{2}\|x\| + \|ux - xu\| \end{aligned}$$

Since $x \in C = \overline{f(u)If(u)} = \overline{h(u)f(u)If(u)h(u)}$ we have:

$$\|ux - xu\| = \|(u - \lambda_0)x - x(u - \lambda_0)\| < \frac{\epsilon}{2}\|x\|$$

Therefore:

$$\|\alpha_{g_0}(x) - x\| < \epsilon\|x\| \quad (1)$$

Let now $\pi \in \Gamma_\omega(\alpha)$. Since $C \in \mathcal{H}^\alpha(A)$, it follows that $\pi \in Sp(\alpha)$. Then $C_\pi \neq (0)$. By Lemma 6 (vii) there is $a \in C$ such that $P_{ii}(\pi)(a) = a_{ii} \neq 0$ for every $i = 1, 2, \dots, d_\pi$.

From relation (1) with $x = a_{ii}$, we have:

$$\|\alpha_{g_0}(a_{ii}) - a_{ii}\| < \epsilon\|a_{ii}\| \quad (2)$$

Lemma 6 (v) and (2) above give:

$$\left\| \sum_{k=1, d_\pi} a_{ik}\pi_{ki}(g_0) - a_{ii} \right\| < \epsilon\|a_{ii}\| \quad (3)$$

where $a_{ik} = P_{ik}(\pi)(a_{ii})$.

By Lemma 6 (vi) and Hahn-Banach there is a functional φ of norm one on C such that $\varphi(a_{ii}) = \|a_{ii}\|$ and $\varphi(a_{ik}) = 0$ for all $k \neq i$.

Applying φ to (3), we get:

$$|\pi_{ii}(g_0) - 1| \cdot \|a_{ii}\| < \epsilon\|a_{ii}\| \quad (4)$$

Hence $|\pi_{ii}(g_0) - 1| < \epsilon$ for every $\epsilon > 0$. Therefore $\pi_{ii}(g_0) = 1$ for all $i = 1, 2, \dots, d_\pi$. Since $\pi(g_0) = [\pi_{ij}(g_0)]$ is a unitary matrix with all diagonal entries equal 1, it follows immediately that $\pi(g_0) = 1$ and we are done. $\square$

Acknowledgements. R. Dumitru was supported by "Isabel and Mary Neff" Fellowship and by URC Summer Grant. C. Peligrad was supported by a Taft Travel Grant. B. Visinescu was supported by a Taft Enhancement Fellowship.

References

- [1] P. Ara and M. Mathieu, Local multipliers of C^* -algebras, Springer 2003
- [2] O. Bratteli, Derivations, dissipations and group actions on C^* -algebras, Springer, Lecture Notes in Math., 1986.
- [3] O. Bratteli, G. A. Elliott, D.E. Evans and A. Kishimoto, Quasi-product actions of a compact abelian group on a C^* -algebra, Tôhoku Math. J. 41(1989), 133-161.
- [4] G. A. Elliott, Automorphisms determined by multipliers on ideals of a C^* -algebra, J. Funct. Anal. 23(1976), 1-10.
- [5] E. C. Gootman, A. J. Lazar and C. Peligrad, Spectra for compact group actions, J. Operator Theory, 31(1994), 381-399.
- [6] D. Olesen, Inner automorphisms of simple C^* -algebras, Comm. Math. Phys. 44(1975), 175-190.
- [7] D. Olesen, G. K. Pedersen and E. Stormer, Compact abelian groups of automorphisms of simple C^* -algebras, Inventiones Math. 39(1977), 55-64.
- [8] G. K. Pedersen, Approximating derivations on ideals of a C^* -algebra, Inventiones Math. 45(1978), 299-305.
- [9] G. K. Pedersen, C^* -algebras and their automorphism groups, Academic Press, London 1979.
- [10] C. Peligrad, Locally compact group actions on C^* -algebras and compact subgroups, J. Funct. Anal. 76(1988), 126-139.
- [11] N. E. Wegge-Olsen, K-theory and C^* -algebras, Oxford University Press 1993.

Costel Peligrad
Department of Mathematics
University of Cincinnati
Cincinnati, Ohio, 45221 – 0025
E – mail : peligrac@math.uc.edu

Raluca Dumitru and Bogdan Visinescu
Department of Mathematics
University of Cincinnati
Cincinnati, Ohio, 45221 – 0025
E – mail : dumitrur@math.uc.edu
E – mail : visinebc@math.uc.edu

On leave from :
Institute of Mathematics
of the Romanian Academy
P.O. Box 1 – 764
70700 Bucharest
Romania